

Advanced Magnetohydrodynamics, by Hans Goedbloed, Rony Keppens and Stefaan Poedts (Cambridge University Press, 2010)

Errata, 19 September 2011

– p. 94, lines after Eq. (13.103):

$\mathbf{H}_A \Rightarrow \mathbf{H}_a$ (thrice)

– p. 108, Fig. 13.14:

There are five (instead of six) unstable eigenvalues, both for the static case (a), and for the rigidly rotating case (b). For case (a), the corrected eigenvalues are $\sigma = 0$, $\nu = 0.3155$, 0.2299 , 0.1724 , 0.1258 and 0.8091 . For case (b), they are $(\sigma, \nu) = (-0.5872, 0.3032)$, $(-0.5580, 0.2225)$, $(-0.5435, 0.1668)$, $(-0.5348, 0.1208)$ and $(-0.5290, 0.7550)$.

– p. 110, Fig. 13.15:

For the value $\lambda = 0.84$, there should be four (instead of three) unstable eigenvalues on the circle, there is no “intruder” mode labelled 1 in that case. If the value of λ is increased to 0.92 , a stable “intruder” mode 1 ($\sigma = -2.102$) appears and two unstable modes 2 ($\sigma = -2.054$, $\nu = 0.0851$) and 3 ($\sigma = -2.000$, $\nu = 0.0633$) appear on the circle. The text on p. 110 then applies without change. The three rightmost eigenvalues become $\sigma \approx -1.263$, -1.116 , -0.785 .

– p. 111, Eq. (13.169):

$< 1 \Rightarrow > 1$

– p. 107–112, Section 13.4.1:

The unstable eigenvalues of the rigidly rotating cylindrical plasma with constant pitch magnetic field are located on the circle given by Eqs. (13.161), (13.165). Contrary to what is frequently stated in the text of Section 13.4.1, this circle is *not* a solution path in the sense of Section 12.3 (which demands $W_2 = 0$), but just a convenient auxiliary curve obtained for the very special case of the Bessel function solutions (13.156). Actually, $W_2 \neq 0$ on that circle.

– p. 116, line after Eq. (13.185):

$r_A \Rightarrow r_a$

– p. 120, 9th line from below:

“over the central part of the imaginary axis” \Rightarrow “over the solution path that slightly (not visibly) deviates [53] from the central part of the imaginary ω axis”

2

– p. 151, Eq. (14.94):

Add a term $-\frac{2r}{m}(n+m/q)(1/q)'$ in the second square bracket, and add “with constant density,” to the line above this equation.

– p. 153, line below Eq. (14.106):

dispersion \Rightarrow dispersion

– p. 288, 3rd line of Eq. (16.164):

RHS of the expression for $\alpha \mathbf{j}_p$ should be multiplied with \mathbf{B}_p .

– p. 293, Eqs. (16.184) and (16.185):

$$z(w) = \frac{w + \delta}{1 + \delta z} \Rightarrow z(w) = \frac{w + \delta}{1 + \delta w}$$
$$\frac{1}{h^2} \left(\frac{1}{s} \frac{\partial}{\partial s} s \frac{\partial \psi}{\partial s} \right) \Rightarrow \frac{1}{h^2} \left(\frac{1}{s} \frac{\partial}{\partial s} s \frac{\partial \psi}{\partial s} + \frac{1}{s^2} \frac{\partial^2 \psi}{\partial t^2} \right)$$

– p. 620, Ref. [331]:

critère \Rightarrow critère